

STAY-C® 50

Exploring the Untapped Potential of
Vitamin C for Skin, Hair and Oral Care

a **DSM** Product



DSM Nutritional Products

STAY-C® 50



STAY-C®

STAY-C® 50, stable Vitamin C

High awareness of vitamins

Consumers think of vitamins as the most important cosmetic ingredients next to UV filters. No other group of ingredients has gained such a high awareness level as vitamins in skin care products. Consumers believe that vitamins deliver many benefits such as wrinkle reduction, firming and fresh appearance¹. This highly positive attitude gives personal care products containing vitamins a head start at the point of sales.

STAY-C® 50 – the stable Vitamin C with numerous benefits for skin, hair and oral care

Vitamin C, under-used for some years due to stability hurdles of straight ascorbic acid, moved into the center of interest with the availability of ascorbic acid derivatives such as the more stable and easy to handle STAY-C® 50.




Figure 1: Structure of STAY-C® 50, sodium salt of the monophosphate ester of ascorbic acid. INCI: sodium ascorbyl phosphate.

Vitamin C is typically referred to as an antioxidant – especially in the combination with Vitamin E – and as an essential substance for collagen synthesis². In addition, several studies showed that STAY-C® 50 is an active principle for a lot more effects not only for skin but also for hair, teeth and gum.

STAY-C® 50



50

STAY-C® 50 and skin

STAY-C® 50 – for visibly younger looking skin

Photo-damage of skin including premature skin aging is to a large extent caused by reactive oxygen species (ROS). Their generation and also the quenching of ROS in living skin can be quantified and visualized. UV filters already protect against UV-induced ROS generation. However this first line of defense is not complete. In vivo antioxidants STAY-C® 50 and Vitamin E acetate were shown to provide an additional quenching effect of ROS of almost 50% to a standard sunscreen formulation of SPF 8 (figure 2)³.




Figure 2: Quenching of UV-induced ROS in human ex vivo skin after the test emulsions have been applied for 3 hours. Dihydrorhodamine was used to monitor the ROS by fluorescence signals. The images, based on two-photon fluorescence microscopy, are taken from the spinous layer of the epidermis. Compared to the SPF 8-control formulation the addition of 2.5% Vitamin E acetate leads to a 23% reduction of the ROS level. The combination of 2.5% STAY-C® 50 and 2.5% Vitamin E acetate quenched the ROS to almost 50%.

STAY-C® 50 does not only reduce the level of damaging free radicals but also helps to increase the skin firmness. Volunteers who had regularly used an emulsion with STAY-C® 50 experienced a firmer skin. This in vivo effect was supported by the additional finding that STAY-C® 50 promotes the synthesis of collagen I and III in human fibroblasts⁴.

STAY-C® 50



STAY-C®

STAY-C® 50 – for better looking skin tone

Vitamin C and some derivatives have been traditionally used as active ingredients to brighten the complexion because they are known to inhibit the formation of melanin. Studies have shown that STAY-C® 50 reduces the melanogenesis in primary human melanocytes by 57%⁵ (figure 3).




Figure 3: A 4-day treatment of primary human melanocytes of skin type II/III with 0.14% STAY-C 50[®] resulted in a 57% reduction of the melanin content. Melanin was quantified by absorption at 475 nm.

Age spots are another visual indication of aged skin resulting from UV light stimulated overproduction of melanin. A 3-month in vivo study on Asian skin has shown that regular use of a cream with STAY-C® 50 resulted in a reduced color intensity of facial age spots⁶. The overall skin lightening effect of STAY-C® 50 was assessed as 25% by dermatological reading (figure 4).




Figure 4: Dermatological assessment of the age spot fading effect of STAY-C® 50 from a 3-months double blind placebo-controlled in vivo study with 39 female volunteers in Asia. The test cream contained 3% STAY-C® 50 and 1% Vitamin E acetate.

The Japanese Ministry of Health has approved STAY-C® 50 as a quasi drug (Q/D) for skin whitening products.

STAY-C® 50

50



STAY-C® 50 – reducing the appearance of acne

Skin blemishes and acne are skin disorders not only experienced by teenagers but also by adults. The emergence of skin impurities and inflammation is complex and involves among many factors the bacterial colonization by *Propionibacterium acnes*. Also the free radical induced peroxidation of sebum is considered to play a role⁷.

Our study using 20 volunteers has shown that an emulsion with 1% STAY-C® 50 significantly inhibits the UVA light induced sebum (squalene) oxidation by 30%. We also could demonstrate that 1% STAY-C® 50 very effectively reduced the bacterial count of *Propionibacterium acnes* under physiological conditions⁵.

Another study

performed together with Professor Ikeno⁸ further illustrates the potential of STAY-C® 50 to improve the appearance of skin conditions associated with pimples and inflammation. Volunteers with moderate to severe skin blemishes regularly applied a 5% STAY-C® 50 lotion twice daily on their face. After 12 weeks their skin condition was significantly ameliorated by 77% on average (figure 5). The visual improvement is illustrated in figure 6 by a representative subject. The efficacy of STAY-C® 50 was superior than 5% Benzoylperoxide, a widely prescribed drug for acne treatment.




Figure 5: Efficacy assessment in 5 grades after 12 weeks twice daily application of either 5% STAY-C® 50 or 5% Benzoylperoxide onto the facial skin of volunteers (2 groups of 30 persons) with inflammatory and non-inflammatory acne lesions.




Figure 6: Representative subject with inflammatory and non-inflammatory lesions of acne vulgaris before treatment (left) and after 12 weeks (right) of twice daily application of a 5% STAY-C® 50 lotion.

STAY-C® 50



STAY-C® 50

STAY-C® 50 – in body care, hair care and oral care

STAY-C® 50 – deodorant activity

Don't be surprised if consumers ask specifically for deodorant products with STAY-C® 50. They might have heard that STAY-C® 50 helps to control body odor. An underarm sniff test involving 10 volunteers revealed that a 0.2% STAY-C® 50 formulation reduced the axillary malodor by more than 40% 8 hours after the application⁴. 24 hours later, the STAY-C® 50 formulation performed an additional 10% better than the placebo.

STAY-C® 50 – hair color protection

Women and men enjoy today's selection of brilliant hair colors and the ease of application. But they also want their color to stay as long as possible. Shampoos and treatment products often contain UV filters to prevent fading or color change. The protective potential of antioxidant vitamins for hair has not been evaluated so far. A study performed on behalf of DSM Nutritional Products showed that the application of a hair tonic with 0.5% STAY-C® 50 onto bleached and permanently colored human hair tresses significantly stabilized the hair color during a 24 hours irradiation time⁴.

STAY-C® 50 – support of healthy teeth and gum

In spite of more regular toothbrushing plaque formation and caries or gum bleeding and gingivitis have remained a problem for many consumers. In both cases, bacteria are strongly involved. Studies have shown that STAY-C® 50 is able to suppress the growth of several bacteria associated with caries or gingivitis. An example of its activity is demonstrated in figure 7. STAY-C® 50 inhibits the growth of *Streptococcus mutans*, the principle causative bacterium of caries. Similar efficacy was also shown against *Actinobacillus actinomycetemcomitans* and *Porphyromas gingivalis*, two bacteria associated with gingivitis⁹.




Figure 7: 0.1 to 1.0% STAY-C® 50 was added to suspensions of *Streptococcus mutans* and incubated for up to 56 hours. An aliquot was plated on colony forming plates, cultivated and statistically counted. STAY-C® 50 inhibited the growth of *S. mutans* at all tested concentrations.

STAY-C® 50

50

Formulating with STAY-C® 50

STAY-C® 50 is a versatile Vitamin C derivative for many applications and product types. From skin care to hair and oral care products it can be included in emulsions, gels, sprays and stick products.

Recommended use level: 0.5 to 5.0%

Some formulation guidelines

- **Maintain the formulation at a pH of around 7 to avoid hydrolysis.**
- **Always use chelating agents (e.g. disodium EDTA) to avoid degradation of STAY-C® 50 by heavy metal ions.**
- **If the formulation is heated, add STAY-C® 50 in the down-cooling step below 40°C. In most cases, the powder can be directly added.**

Different reference formulations are available and can be sent on request.

References

- 1 Consumer Market Research Study by GfK, sponsored by DSM Nutritional Products (2000)
- 2 N Boyera, I Galey, BA Bernard, Effect of Vitamin C and its derivatives on collagen synthesis and cross-linking by normal human fibroblasts. *Int J Cosm Sci* 20 151-158 (1998)
- 3 KM Hanson & RM Clegg, Bioconvertible vitamin antioxidants improve sunscreen protection against UV-induced reactive oxygen species. *Journal of Cosmetic Science* 54 (6) 589-598 (2003)
- 4 R Jermann & AA Gripp, STAY-C® 50 – more than just stable Vitamin C in your formulation. *Cosmetics and Toiletries Manufacture Worldwide* 153-157 (2004)
- 5 J Klock, Sodium Ascorbyl Phosphate has still unexploited potential in cosmetics. *SÖFW* 130 46-56 (2004)
- 6 Asia-Pacific Skin Testing Center, Facial cream 12-week controlled usage study, sponsored by DSM Nutritional Products (2002)
- 7 D Saint-Leger, A Bague, E Cohen, M Chivot, A possible role for squalene in the pathogenesis of acne. I. In vitro study of squalene oxidation. *Br J Dermatol* 114 535-542 (1986)
- 8 J Klock, H Ikeno, K Ohmori, J Vollhardt, V Schehlmann, Sodium ascorbyl phosphate shows in vitro and in vivo efficacy in the prevention and treatment of acne vulgaris. In: *Proceedings of the IFSCC-Congress, Orlando Florida, 24-26 October (2004)*
- 9 D Walther, Mikrobiologischer Standzeittest, sponsored by DSM Nutritional Products (2002)

STAY-C® 50



North America

DSM Nutritional Products, Inc.
45 Waterview Blvd.
Parsippany, NJ 07054-1298
U.S.A.
Telephone +1 800 526 0189
Telefax +1 973 257 8420
Internet <http://www.nutraaccess.com>

Asia

DSM Nutritional Products Asia Pacific Pte, Ltd.
78 Shenton Way, # 15-02 LIPPO Center
Singapore 079120
Singapore
Telephone +65 6 325 6200
Telefax +65 6 220 1976
E-mail: unlimited.personalcare@dsm.com

Latin America

DSM Nutritional Products Latin America
Av. Engenheiro Billings 1729 – Predio 9
05321-900 – São Paulo – SP
Brazil
Telephone +55 11 3719 4604
Telefax +55 11 3719 4985
Internet <http://www.dsmnutritionalproducts.com>
E-mail: américa-latina.dnp@dsm.com

Europe

DSM Nutritional Products Europe
P.O. Box 3255
CH-4002 Basel
Switzerland
Telephone +41 61 687 17 77
Telefax +41 61 688 35 89
Internet <http://www.dsmnutritionalproducts.com>

The information given in this publication is based on our current knowledge and experience, and may be used at your discretion and risk. It does not relieve you from carrying out your own precautions and tests. For some countries there is mention of uses beyond the scope of the approved or accepted labeling for the subject product. We do not assume any liability in connection with your product or its use. You must comply with all applicable laws and regulations, and observe all third party rights.

STAY-C® 50